

ON EXTREMAL CONTRACTIONS FROM THREEFOLDS TO SURFACES: THE CASE OF ONE NON-GORENSTEIN POINT

YURI G. PROKHOROV

In Memoriam: Professor Wei-Liang Chow

ABSTRACT. Let $f: X \rightarrow S$ be an extremal contraction from a threefold with only terminal singularities to a surface. We study local analytic structure such contractions near degenerate fiber C in the case when C is irreducible and X has on C only one non-Gorenstein point.

INTRODUCTION

Let V be a non-singular algebraic projective threefold over $\mathbb{C}$ of Kodaira dimension $\kappa(V) = -\infty$. It follows from the Minimal Model Program that V is birationally equivalent to an algebraic projective threefold X with only terminal $\mathbb{Q}$ -factorial singularities and with a fiber space structure $f: X \rightarrow S$ over a lower dimensional variety S such that $-K_X$ is f -ample and relative Picard number of X/S is one. Thus, at least in principle, one can reduce the problem of studying birational structure of V to studying structure of S and the fiber space X/S . In the present paper we investigate the local structure of X/S near singular fibers in the case $\dim(S) = 2$. In this situation it is easy to see that a general fiber is a conic in $\mathbb{P}^2$. We will work in the analytic situation.

Definition 0.1. Let (X, C) be a germ of a three-dimensional complex space along a compact reduced curve C and let $(S, 0)$ be a germ of a two-dimensional normal complex space. Suppose that X has at worst terminal singularities. Then we say that a proper morphism $f: (X, C) \rightarrow (S, 0)$ is a *Mori conic bundle* if

- (i) $(f^{-1}(0))_{\text{red}} = C$;
- (ii) $f_*\mathcal{O}_X = \mathcal{O}_S$;
- (iii) $-K_X$ is ample.

The following conjecture is interesting for applications of Sarkisov program to the rationality problem for conic bundles [6], [5] or studying of $\mathbb{Q}$ -Fano threefolds with extremal contractions onto a surface [17].

1991 *Mathematics Subject Classification*. Primary 14E30; Secondary 14E05, 14E35, 14J30.

The author was supported in part by the Russian Foundation of Fundamental Research # 96-01-00820 and the International Science Foundation # M9O 300.

Conjecture 0.2. *Let $f: (X, C) \rightarrow (S, 0)$ be a Mori conic bundle. Then $(S, 0)$ is a DuVal singularity of type A_n .*

It was proved in [3] that if X is Gorenstein, then $(S, 0)$ is non-singular and $f: (X, C) \rightarrow (S, 0)$ is a "usual" conic bundle, i. e. there exists an embedding $(X, C) \subset \mathbb{P}^2 \times (S, 0)$ such that all the fibers of f are conics in $\mathbb{P}^2$. Earlier [18] (see also [19]) we classified all Mori conic bundles containing only points of indices ≤ 2 . In this case $(S, 0)$ is non-singular or DuVal of type A_1 . In general, Conjecture 0.2 follows from the following special case of Reid's general elephant conjecture (see [16], [17]).

Conjecture 0.3. *Let $f: (X, C) \rightarrow (S, 0)$ be a Mori conic bundle. Then the linear system $| -K_X |$ contains a divisor having only DuVal singularities.*

It can be generalized as follows.

Conjecture 0.4. *Let $f: (X, C) \rightarrow (S, 0)$ be a Mori conic bundle. Then for any $n \in \mathbb{N}$ the following holds*

$(*_n)$ *There exists a divisor $D \in | -nK_X |$ such that $K_X + \frac{1}{n}D$ is log-terminal.*

(Note that 0.4 $(*_1)$ is equivalent to 0.3).

By [7, Theorem 4.5], for every threefold X with terminal singularities there exists a projective bimeromorphic morphism $q: X^q \rightarrow X$ called $\mathbb{Q}$ -factorialization of X such that X^q has only terminal $\mathbb{Q}$ -factorial along $q^{-1}(C)$ singularities and q is an isomorphism in codimension 1. If $f: (X, C) \rightarrow (S, 0)$ is a Mori conic bundle, then applying the Minimal Model Program to X^q over $(S, 0)$ we obtain a new Mori conic bundle $f': (X', C') \rightarrow (S, 0)$ over the same base surface S and $\rho(X', C')/(S, 0)=1$. In particular C' is irreducible by Corollary 1.1.1. Therefore it is sufficient to prove Conjecture 0.2 assuming that C is irreducible.

In the this paper we will study Mori conic bundles $f: (X, C) \rightarrow (S, 0)$ with irreducible central fiber. In this situation X contains at most three singular points (Corollary 2.7). Arguments similar to [14, 0.4.13.3] show also that X can contain at most two non-Gorenstein points [17]. Our results concern only the case when X contains only one non-Gorenstein singular point. It will be proved in this case (Theorem 4.1) that the point $(S, 0)$ may be either non-singular or DuVal of type A_1 or A_3 . We also show that either $(*_1)$ or $(*_2)$ hold and give a classification if $(S, 0)$ is singular. Our main tool will be Mori's technique [14].

Acknowledgments. I would like to acknowledge discussions that I had on this subject with Professors V.I. Iskovskikh, V.V. Shokurov and M. Reid. I have been working on this problem at Max-Planck Institut für Mathematik and the Johns Hopkins University. I am very grateful them for hospitality and support.

1. PRELIMINARY RESULTS

The following is a consequence of the Kawamata-Viehweg vanishing theorem (see [15, §4], [8, 1-2-5]).

Theorem 1.1. *Let $f: (X, C) \rightarrow (S, 0)$ be a Mori conic bundle. Then $R^i f_* \mathcal{O}_X = 0$, $i > 0$.*

□

Corollary 1.1.1. (cf. [14, (1.2)-(1.3)]) *Let $f: (X, C) \rightarrow (S, 0)$ be a Mori conic bundle. Then*

- (i) *For an arbitrary ideal $\mathcal{I}$ such that $\text{Supp}(\mathcal{O}_X/\mathcal{I}) \subset C$ we have, $H^1(\mathcal{O}_X/\mathcal{I}) = 0$.*
- (ii) *The fiber C is a tree of non-singular rational curves (i. e. $p_a(C') \leq 0$ for any subcurve $C' \subset C$).*
- (iii) *If C has ρ irreducible components, then*

$$\text{Pic}(X) \simeq H^2(C, \mathbb{Z}) \simeq \mathbb{Z}^{\oplus \rho}.$$

□

1.2. Terminal singularities. Let (X, P) be a terminal singularity of index $m \geq 1$ and let $\pi: (X^\sharp, P^\sharp) \rightarrow (X, P)$ be the canonical cover. Then $(X^\sharp, P^\sharp)$ is a terminal singularity of index 1 and $(X, P) \simeq (X^\sharp, P^\sharp)/\mathbb{Z}_m$. It is known (see [20]) that $(X^\sharp, P^\sharp)$ is a hypersurface singularity, i. e. there exist an $\mathbb{Z}_m$ -equivariant embedding $(X^\sharp, P^\sharp) \subset (\mathbb{C}^4, 0)$. Fix a character χ that generates $\text{Hom}(\mathbb{Z}_m, \mathbb{C}^*)$. For $\mathbb{Z}_m$ -semi-invariant z define weight $\text{wt}(z)$ as an integer defined mod m such that

$$\text{wt}(z) \equiv a \pmod{m} \quad \text{iff} \quad \zeta(z) = \chi(\zeta)^a \cdot z \quad \text{for all} \quad \zeta \in \mathbb{Z}_m.$$

Theorem 1.2.1. (see [13], [21]) *In notations of 1.2 $(X^\sharp, P^\sharp)$ is analytically $\mathbb{Z}_m$ -isomorphic to a hypersurface $\phi = 0$ in $(\mathbb{C}_{x_1, x_2, x_3, x_4}^4, 0)$ such that x_1, x_2, x_3, x_4 are semi-invariants and we have one of the following two series:*

- (i) *(the main series) $\text{wt}(x_1, x_2, x_3, x_4; \phi) \equiv (a, -a, b, 0; 0) \pmod{m}$, or*
- (ii) *(the exceptional case) $m = 4$ and $\text{wt}(x_1, x_2, x_3, x_4; \phi) \equiv (a, -a, b, 2; 2) \pmod{4}$,*

where a, b are integers prime to m .

Note that in case (i) the variety $X^\sharp$ can be non-singular. In this situation $(X, P) \simeq (\mathbb{C}^3, 0)/\mathbb{Z}_m(a, -a, b)$. Such singularities are called *terminal cyclic quotient singularities* and denoted also by $\frac{1}{m}(a, -a, b)$.

Theorem 1.2.2. (see [21]) *Let (X, P) be a germ of terminal singularity. Then a general member $F \in |-K_X|$ has only DuVal singularity (at P).*

1.2.3. Terminal singularities can be classified in terms of a general member $F \in |-K_X|$ and its preimage $F^\sharp := \pi^{-1}(F)$ under the canonical cover $\pi: X^\sharp \rightarrow X$ [21]:

index	type	cover $F^\sharp \rightarrow F$
the main series		
m	cA/m	$A_{k-1} \xrightarrow{m:1} A_{km-1}$
2	$cAx/2$	$A_{2k-1} \xrightarrow{2:1} D_{k+2}$
2	$cD/2$	$D_{k+1} \xrightarrow{2:1} D_{2k}$
3	$cD/3$	$D_4 \xrightarrow{3:1} E_6$
2	$cE/2$	$E_6 \xrightarrow{2:1} E_7$
the exceptional case		
4	$cAx/4$	$A_{2k-2} \xrightarrow{4:1} D_{2k+1}$

Cyclic quotient singularities are included in type cA/m .

Definition 1.2.4. Let X be a normal variety and $\text{Cl}(X)$ be its Weil divisor class group. The subgroup of $\text{Cl}(X)$ consisting of Weil divisor classes which are $\mathbb{Q}$ -Cartier is called the *semi-Cartier divisor class group*. We denote it by $\text{Cl}^{\text{sc}}(X)$.

Theorem 1.2.5. (see [20],[7]) *Let (X, P) be a germ of 3-dimensional singularity. Then $\text{Cl}^{\text{sc}}(X, P) \simeq \mathbb{Z}_m$ and it is generated by the class of K_X . Moreover the local fundamental group of (X, P) is cyclic: $\pi_1(X - \{P\}) \simeq \mathbb{Z}_m$.*

Definition 1.2.6. Let $(S, 0)$ be a two-dimensional log-terminal singularity. It is well known (see [7]) that $(S, 0)$ is a quotient singularity, i. e. $(S, 0) \simeq (\mathbb{C}^2, 0)/G$, where $G \subset GL(2, \mathbb{C})$ is a finite subgroup without quasi-reflections. The natural cover $(\mathbb{C}^2, 0) \rightarrow (S, 0)$ we call the *topological cover* and the order of G we call the *topological index* of $(S, 0)$.

The following is an easy consequence of 1.2.

Lemma 1.2.7. *Let (X, P) be a germ of a terminal threefold singularity of index $m > 1$ and $(F, P) \subset (X, P)$ be a germ of irreducible surface. Assume that F is $\mathbb{Q}$ -Cartier and (F, P) is DuVal with topological index n .*

- (i) *Then n is divisible by m .*
- (ii) *Moreover if $n = m$, then (X, P) is a cyclic quotient singularity and (F, P) is of type A_{m-1} .*

Proof. Let $F^\sharp := \pi^{-1}(F)$. The divisor $F^\sharp$ is Cartier on $X^\sharp$, because point $(X^\sharp, P^\sharp)$ is terminal of index 1 (see [7, Lemma 5.1]). On the other hand $F^\sharp$ is non-singular outside $P^\sharp$. Therefore $F^\sharp$ is irreducible and normal. The cover $F^\sharp \rightarrow F$ is étale outside P . Hence the topological cover of (F, P) can be factorized through $F^\sharp \rightarrow F$. This proves (i). In conditions (ii) $F^\sharp$ is non-singular and so is $X^\sharp$. $\square$

Proposition 1.3. (see [14]) Let (X, P) be a germ of terminal singularity of index m , $(C, P) \subset (X, P)$ be a germ of smooth curve and $\pi: (X^\sharp, P^\sharp) \rightarrow (X, P)$ be the canonical cover and $C^\sharp := (\pi^{-1}(C))_{\text{red}}$. Then

- (i) for an arbitrary $\xi \in \text{Cl}^{sc}(X, P)$, there exists an effective (Weil) divisor D such that $[D] = \xi$ and $\text{Supp}(D) \cap C = \{P\}$.
- (ii) $\xi \rightarrow (D \cdot C)_P$ induces a homomorphism

$$\text{cl}(C, P): \text{Cl}^{sc}(X, P) \rightarrow \frac{1}{m}\mathbb{Z}/\mathbb{Z} \subset \mathbb{Q}/\mathbb{Z}.$$

□

Definition 1.4. (see [14]) Let things be as in 1.3. $X \supset C$ is called (locally) *imprimitive* at P of *splitting degree* d if $\text{cl}(C, P): \text{Cl}^{sc}(X, P) \rightarrow \frac{1}{m}\mathbb{Z}/\mathbb{Z}$ is not an isomorphism and the order of $\text{Ker}(\text{cl}(C, P))$ is equal d . The order of $\text{Coker}(\text{cl}(C, P))$ is called the *subindex* of P and usually denoted by $\bar{m}$. (Note that $\bar{m}d = m$). If $\text{cl}(C, P): \text{Cl}^{sc}(X, P) \rightarrow \frac{1}{m}\mathbb{Z}/\mathbb{Z}$ is an isomorphism, then P is said to be (locally) *primitive*. In this case we put $d = 1$, $\bar{m} = m$.

In the situation above the preimage $C^\sharp := \pi^{-1}(C)$ under the canonical cover $\pi: X^\sharp \rightarrow X$ has exactly d irreducible components.

1.5. Construction, [14]. Let $f: (X, C \simeq \mathbb{P}^1) \rightarrow (S, 0)$ be a Mori conic bundle with irreducible central fiber and let $P \in X$ be a point of index $m > 1$. Assume that (X, C) is imprimitive of splitting degree d at P . Take an effective Cartier divisor H such that $H \cap C$ is a smooth point of X and $(H \cdot C) = 1$ and take an effective Weil $\mathbb{Q}$ -Cartier divisor D such that $D \cap C = \{P\}$ and D is a generator of $\text{Cl}^{sc}(X, P)$ (see 1.3). Then the divisor

$$D := (m/d)D - ((mD \cdot C)/d)H$$

is a d -torsion element in $\text{Cl}^{sc}(X, C)$. It defines a finite Galois $\mathbb{Z}_d$ -morphism $g^b: X' \rightarrow X$ such that $P' := g'^{-1}(P)$ is one point, g' is étale over $X - \{P\}$ (hence X' has only terminal singularities), index of (X', P') is equal to m/d , $C' := (g'^{-1}(C))_{\text{red}}$ is a union of d $\mathbb{P}^1$'s meeting only at P' , and each irreducible component of C' is primitive at P' .

2. LOCAL INVARIANTS w_P AND i_P

In this section we following Mori [14] introduce numerical invariants w_P and i_P . There is nothing new in this section. The material is contained only for convenience of the reader.

2.1. Let X be a normal three-dimensional complex space with only terminal singularities and let $C \subset X$ be a reduced non-singular curve. Denote by $\mathcal{I}_C$ the ideal sheaf of C and $\omega_X := \mathcal{O}_X(K_X)$. As in [14], we consider the following sheaves on C :

$$\text{gr}_C^0 \omega := \text{torsion-free part of } \omega_X / (\mathcal{I}_C \omega_X),$$

$$\text{gr}_C^1 \mathcal{O} := \text{torsion-free part of } \mathcal{I}_C / \mathcal{I}_C^2.$$

Since C is non-singular, we have

$$\omega_X/(\mathcal{I}_C\omega_X) = \mathrm{gr}_C^0\omega \oplus \mathrm{Tors}, \quad \mathcal{I}_C/\mathcal{I}_C^2 = \mathrm{gr}_C^1\mathcal{O} \oplus \mathrm{Tors}.$$

2.2. Let m be the index of X . The natural map

$$(\omega_X \otimes \mathcal{O}_C)^{\otimes m} \rightarrow \mathcal{O}_C(mK_X)$$

induces an injection

$$\beta: (\mathrm{gr}_C^0\omega)^{\otimes m} \rightarrow \mathcal{O}_C(mK_X).$$

Denote

$$w_P := (\mathrm{length}_P \mathrm{Coker} \beta)/m.$$

Note that if $f: (X, C \simeq \mathbb{P}^1) \rightarrow (S, 0)$ is a Mori conic bundle, then $\deg \mathrm{gr}_C^0\omega < 0$, (because $\deg \mathcal{O}_C(mK_X) < 0$).

2.3. We also have the natural map

$$\begin{aligned} \mathrm{gr}_C^1\mathcal{O} \times \mathrm{gr}_C^1\mathcal{O} \times \omega_C &\longrightarrow \omega_X \otimes \mathcal{O}_C \rightarrow \mathrm{gr}_C^0\omega, \\ x \times y \times zdt &\longrightarrow zdx \wedge dy \wedge dt \end{aligned}$$

that induces a map

$$\alpha: \wedge^2(\mathrm{gr}_C^1\mathcal{O}) \otimes \omega_C \rightarrow \mathrm{gr}_C^0\omega.$$

Let

$$i_P := \mathrm{length}_P \mathrm{Coker}(\alpha).$$

Note that $i_P = 0$ if X is smooth at P .

Lemma 2.3.1. ([14, 2.15]) *If (X, P) is singular, then $i_P \geq 1$.*

□

Example 2.4. Let (X, P) be a cyclic quotient of type $1/\bar{m}(1, -1, \bar{m}+1)$ ($\bar{m}$ is even) and let $\pi: (\mathbb{C}_{x_1, x_2, x_3}^3, 0) \rightarrow (X, P)$ be its canonical cover. Consider the curve $C^\# := \{x_3 = x_2^2 - x_1^{2\bar{m}-2} = 0\} \subset \mathbb{C}^3$. This curve has exactly two components $C^\#(1), C^\#(2) = \{x_3 = x_2 \pm x_1^{\bar{m}-1} = 0\}$ permuted under the action of $\mathbb{Z}_{2\bar{m}}$. Thus the image $C := \pi(C^\#)$ is a smooth irreducible curve, X is locally imprimitive along C with splitting degree 2. The curve C is naturally isomorphic to $C^\#(1)/\mathbb{Z}_{\bar{m}}$ and a local uniformizing parameter on C is $x_1^{\bar{m}}$. It is easy to see that

- (i) $\mathcal{O}_{C,P} \simeq \mathbb{C}\{x_1^{\bar{m}}\}$.
- (ii) $\mathcal{O}_C(mK_X) \simeq \mathcal{O}_C(dx_1 \wedge dx_2 \wedge dx_3)^m$, $\mathrm{gr}_C^0\omega \simeq \mathcal{O}_C(x_1^{\bar{m}-1}x_1 \wedge dx_2 \wedge dx_3)^m$.
- (iii) $\mathrm{gr}_C^1\mathcal{O} \simeq \mathcal{O}_C(x_1^{\bar{m}-1}x_3) \oplus \mathcal{O}_C(x_1^2(x_2^2 - x_1^{2\bar{m}-2}))$.
- (iv) $w_P = (\bar{m} - 1)/\bar{m}$, $i_P = 2$.

From definitions we have

Proposition 2.5. *If $C \simeq \mathbb{P}^1$, then*

$$\deg \operatorname{gr}_C^1 \mathcal{O} = 2 + \deg \operatorname{gr}_C^0 \omega - \sum_P i_P,$$

$$(K_X \cdot C) = \deg \operatorname{gr}_C^0 \omega + \sum_P w_P.$$

□

Proposition 2.6. *Let $f: (X, C \simeq \mathbb{P}^1) \rightarrow (S, 0)$ be a Mori conic bundle. Then*

$$\deg \operatorname{gr}_C^1 \mathcal{O} \geq -2.$$

Proof. Consider the exact sequence

$$0 \rightarrow \mathcal{I}_C / \mathcal{I}_C^2 \rightarrow \mathcal{O}_X / \mathcal{I}_C^2 \rightarrow \mathcal{O}_C \rightarrow 0.$$

By Corollary 1.1.1 $H^1(\mathcal{O}_X / \mathcal{I}_C^2) = 0$ and since $H^0(\mathcal{O}_X / \mathcal{I}_C^2) \rightarrow H^0(\mathcal{O}_C)$ is onto, we have $H^1(\mathcal{I}_C / \mathcal{I}_C^2) = 0$. Hence $H^1(\operatorname{gr}_C^1 \mathcal{O}) = 0$. It gives us our assertion. □

Corollary 2.6.1. *Let $f: (X, C \simeq \mathbb{P}^1) \rightarrow (S, 0)$ be a Mori conic bundle. Then*

$$(-K_X \cdot C) + \sum w_P + \sum i_P \leq 4.$$

Remark. In the case of flipping contraction we always have $\deg \operatorname{gr}_C^0 \omega = -1$ (see [14]) but in our situation of Mori conic bundles there are examples (e. g. 4.1.2 (i)) with $\deg \operatorname{gr}_C^0 \omega = -2$. Nevertheless it is easy to see from 2.5 and 2.6 that $-1 \geq \deg \operatorname{gr}_C^0 \omega \geq -3$. J. Kollár observe also that from vanishing $R^1 f_* \mathcal{O}_X(K_X + H) = 0$ for any ample Cartier divisor H , we have $\deg \operatorname{gr}_C^0 \omega = -1$ or -2 .

Corollary 2.7. *Let $f: (X, C \simeq \mathbb{P}^1) \rightarrow (S, 0)$ be a Mori conic bundle. Then (X, C) contains at most three singular points.*

Proof. It follows from Corollary 2.6.1 and Lemma 2.3.1. □

As in [14] we need the following construction to compute invariants w_P and i_P locally.

2.8. Let (X, P) be a terminal point of index m and let $C \subset (X, P)$ be a smooth curve. Let $\pi: (X^\sharp, P^\sharp) \rightarrow (X, P)$ be the canonical cover and $C^\sharp := (\pi^{-1}(C))_{\text{red}}$. We suppose that (X, P) has splitting degree d along C and subindex $\bar{m}$ (i. e. $C^\sharp$ has exactly d components and $m = \bar{m}d$). By 1.2, there exists an $\mathbb{Z}_m$ -equivariant embedding $X^\sharp \subset \mathbb{C}^4$. Let $\phi = 0$ be an equation of $X^\sharp$ in $\mathbb{C}^4$.

Let $C_1^\sharp$ be an irreducible component of $C^\sharp$. For $z \in \mathcal{O}_{X^\sharp}$ we define $\operatorname{ord}(z)$ as the order of vanishing of z on the normalization of $C_1^\sharp$. It is easy to see that $\operatorname{ord}(z)$ does not depend on our choice of a component $C_1^\sharp$, if z is a

semi-invariant. All the values $\text{ord}(z)$ form a semigroup which is denoted by $\text{ord}(C^\sharp)$. Obviously the "coordinate" values $\text{ord}(x_1), \dots, \text{ord}(x_4)$ generate $\text{ord}(C^\sharp)$.

Proposition-Definition 2.8.1. By [14], one can choose a coordinate system x_1, x_2, x_3, x_4 in $\mathbb{C}^4$ and a character $\chi: \mathbb{Z}_m \rightarrow \mathbb{C}^*$ such that the following conditions hold.

- (i) Coordinates x_1, x_2, x_3, x_4 are semi-invariants with weights satisfying 1.2.1.
- (ii) $\text{ord}(x_i) \equiv \text{wt}(x_i) \pmod{\bar{m}}$ for all $i = 1, 2, 3, 4$.
- (iii) Each component of $C^\sharp$ is parameterized as

$$x_i = \chi(g)^{\text{wt}(x_i)} t^{\text{ord}(x_i)}, \quad i = 1, 2, 3, 4, \quad g \in \mathbb{Z}_m / \mathbb{Z}_{\bar{m}}.$$

- (iv) *Normalizedness property.* For any $i = 1, 2, 3, 4$ there is no semi-invariants y such that $\text{wt}(y) \equiv \text{wt}(x_i) \pmod{m}$ and $\text{ord}(y) < \text{ord}(x_i)$. In particular $\text{ord}(x_i) < \infty$ for all i .
- (v) There exists an invariant function z on $X^\sharp$ such that $\text{ord}(z) = \bar{m}$. In particular in the case of the main series one has $\text{ord}(x_4) = \bar{m}$, $\text{wt}(x_4) \equiv 0 \pmod{m}$.

Such a coordinate system is said to be *normalized*. Note that we still may permute x_1, x_2 and may replace a character χ with χ' if $\chi' \equiv \chi \pmod{\bar{m}}$.

□

Proposition 2.9. (see [14, 2.10]) *Under the notations and conditions 2.8.1 one has*

$$w_P = \min\{\text{ord}(\psi) \mid \psi \in \mathcal{O}_{X^\sharp, P^\sharp}, \quad \text{wt}(\psi) \equiv -\text{wt}(x_3) \pmod{m}\}.$$

□

Proposition 2.10. (cf. [14, 0.4.14.2]) *Let $F \in |-K_{(X,P)}|$ be a general member. Then in a normalized coordinate system we have $(F \cdot C)_P = \text{ord}(x_3)/\bar{m}$.*

□

Proposition 2.11. (see [14, 2.12]) *Under the notations of 2.8.1, one has*

$$i_P \bar{m} = \bar{m} - \text{ord}(x_4) - \bar{m} w_P + \min_{\phi_1, \phi_2 \in J_0^\sharp} [\phi, \phi_1, \phi_2],$$

where $J_0^\sharp$ is the invariant part of the ideal sheaf $J^\sharp$ of $C^\sharp$ in $\mathbb{C}^4$, ϕ is an equation of $X^\sharp$ in $\mathbb{C}^4$ and

$$[\psi_1, \psi_2, \psi_3] := \text{ord} \partial(\psi_1, \psi_2, \psi_3) / \partial(x_1, x_2, x_3).$$

is the Jacobian determinant.

□

By semi-additivity of $[\ , \ , \]$ and because $x_4 \phi \in J_0^\sharp$, we have

Corollary 2.11.1. (see [14, (2.15.1)]) *In notations above*

$$i_P \bar{m} \geq -\bar{m} w_P + \min_{\psi_1, \psi_2, \psi_3 \in J_0^\sharp} [\phi_1, \phi_2, \phi_3].$$

□

2.11.2. It is clear that the ideal $J_0^\sharp$ is generated by invariants of the form $\psi - x_4^n$ where ψ is an invariant monomial with $\text{ord}(\psi) = \text{ord}(x_4^n) = n\bar{m}$. Thus we can assume in 2.11.1 that $\phi_i = \psi_i - x_4^{n_i}$, for $i = 1, 2, 3$, where each ψ_i is an invariant monomial in x_1, x_2, x_3 . For any element $\psi = x_1^{b_1} x_2^{b_2} x_3^{b_3} x_4^{b_4} - x_4^n$ define the vector $\text{ex}_4(\psi) \in \mathbb{Z}^3$ by $\text{ex}_4(\psi) = (b_1, b_2, b_3)$.

Lemma 2.11.3. (see [14, 2.14]) *Let ψ_1, ψ_2, ψ_3 be as in 2.11.2. Then*

$$[\psi_1, \psi_2, \psi_3] = \begin{cases} \sum_{i=1}^3 \text{ord}(\psi_i) - \sum_{i=1}^3 \text{ord}(x_i) & \text{if } \text{ex}_4(\psi_i), i = 1, 2, 3 \\ & \text{are independent,} \\ \infty & \text{otherwise.} \end{cases}$$

□

An invariant monomial ψ in x_1, x_2, x_3 is said to be *simple* if it cannot be presented as a product of two non-constant invariant monomials.

Corollary 2.11.4. (see [14, proof of 2.15]) *Under the notations and conditions of 2.8.1 for some simple different invariant monomials ψ_1, ψ_2, ψ_3 in x_1, x_2, x_3 we have*

$$(*) \quad \bar{m} i_P \geq (\text{ord}(\psi_1) - \text{ord}(x_1)) + (\text{ord}(\psi_2) - \text{ord}(x_2)) + (\text{ord}(\psi_3) - \text{ord}(x_3) - \bar{m} w_P).$$

Moreover up to permutations ψ_1, ψ_2, ψ_3 we may assume that $\psi_i = x_i \nu_i$, where $\nu_i, i = 1, 2, 3$ also are monomials. In particular, all three terms in the formula are non-negative and

$$(**) \quad \bar{m} i_P \geq \text{ord}(\nu_1) + \text{ord}(\nu_2) + \text{ord}(\nu_3) - \bar{m} w_P.$$

□

3. EASY LEMMAS

3.1. **Construction.** Let $f: (X, C) \rightarrow (S, 0)$ be a Mori conic bundle. Assume that $(S, 0)$ is singular. Then $(S, 0)$ is a log-terminal point (see e. g. [11] or [4]) and the topological cover $h: (S^b, 0) \simeq (\mathbb{C}^2, 0) \rightarrow (S, 0)$ is non-trivial. Let X^b be the normalization of $X \times_S S^b$ and $G = \text{Gal}(S^b/S)$. Then we have the diagram

$$\begin{array}{ccc} X^b & \xrightarrow{g} & X \\ f^b \downarrow & & \downarrow f \\ S^b & \xrightarrow{h} & S \end{array}$$

The group G acts on X^b and clearly $X = X^b/G$. Since the action of G on $S^b - \{0\}$ is free, so is the action of G on $X^b - C^b$, where $C^b := (f^{b^{-1}}(0))_{\text{red}}$.

Therefore $X^\flat$ has only terminal singularities and the induced action of G on $X^\flat$ is free outside of a finite set of points (see e. g. [2, 6.7]). Since $K_{X^\flat} = g^*(K_X)$, $f^\flat: (X^\flat, C^\flat) \rightarrow (S^\flat, 0)$ is a Mori conic bundle with non-singular base surface.

As an easy consequence of the construction above and 1.2.5 we have

Proposition 3.2. (see [16], [17], [10]) *Let $f: (X, C) \rightarrow (S, 0)$ be a Mori conic bundle. Then $(S, 0)$ is a cyclic quotient singularity.*

Corollary 3.3. *Let $f: (X, C) \rightarrow (S, 0)$ be a Mori conic bundle and let T be the torsion part of the Weil divisor class group $\text{Cl}(X)$. Then*

- (i) *T is a cyclic group,*
- (ii) *the topological index of $(S, 0)$ is equal to the order of T .*

In particular, $(S, 0)$ is non-singular iff $T = 0$.

Proof. Consider the diagram from 3.1. We claim that $\text{Cl}(X^\flat)$ is torsion-free. Indeed in the opposite case there is a finite étale in codimension 1 cover $X' \rightarrow X^\flat$ and then it must be étale outside finite number singular points on $C^\flat$. One can take the Stein factorization $X' \rightarrow S' \rightarrow S^\flat$. Then $S' \rightarrow S^\flat$ will be étale outside 0. This contradicts smoothness of $S^\flat$. Therefore $\text{Cl}(X^\flat)$ is torsion-free and $X^\flat \rightarrow X$ is the maximal abelian cover of X . It gives us $T \simeq \mathbb{Z}_n$, where n is the topological index of $(S, 0)$. $\square$

Proposition 3.4. *Let $f: (X, C \simeq \mathbb{P}^1) \rightarrow (S, 0)$ be a Mori conic bundle with irreducible central fiber. Suppose that the point $(S, 0)$ is singular of topological index $n > 1$. Then for $f^\flat: (X^\flat, C^\flat) \rightarrow (S^\flat, 0)$ of 3.1 we have the following two possibilities.*

- (i) (primitive case) *$C^\flat$ is irreducible, X has at least two non-Gorenstein points.*
- (ii) (imprimitive case) *$C^\flat$ has $d > 1$ irreducible components, they all pass through one point $P^\flat$ and they do not intersect elsewhere. If in this case $m^\flat$ is the index of $P^\flat$ and n is the topological index of $(S, 0)$, then the index of the point $g(P) \in X$ is equal to $nm^\flat$.*

Proof. If $C^\flat$ is irreducible, then the action of $G \simeq \mathbb{Z}_n$ on $C^\flat \simeq \mathbb{P}^1$ has exactly two fixed points $P_1^\flat$ and $P_2^\flat$. Their images $g(P_1^\flat)$ and $g(P_2^\flat)$ are points of indices > 1 . Assume that $C^\flat$ is reducible. Then $G \simeq \mathbb{Z}_n$ acts on components $\{C_i^\flat\}$ of $C^\flat$ transitively. Since $\cup C_i^\flat$ is a tree, $\cap C_i^\flat$ is a point and this point must be fixed under G -action. The rest is obvious. $\square$

Corollary 3.5. *Let $f: (X, C \simeq \mathbb{P}^1) \rightarrow (S, 0)$ be a Mori conic bundle with irreducible central fiber. Assume that X contains only one non-Gorenstein point P . Then the following conditions are equivalent*

- (i) *$(S, 0)$ is singular of topological index n ,*
- (ii) *C is locally imprimitive (see 1.4) at P of splitting degree n .*

Lemma 3.6. *Let $f: (X, C \simeq \mathbb{P}^1) \rightarrow (S, 0)$ be a minimal Mori conic bundle. Assume that f is imprimitive of splitting degree d . Then d is even.*

Proof. Consider the base change 3.1. Then $f^b: (X^b, C^b) \rightarrow (S^b, 0)$ is a Mori conic bundle with $\rho(X^b) = d$. Since $-K_{X^b}$ is f^b -ample, the Mori cone $\overline{NE}((X^b, C^b)/(S^b, 0)) \subset \mathbb{R}^\rho$ is generated by classes of $C_1^b, \dots, C_d^b$. Thus each C_i^b generates an extremal ray R_i and $\overline{NE}((X^b, C^b)/(S^b, 0)) \subset \mathbb{R}^\rho$ is simplicial. The contraction of each extremal face of $\overline{NE}((X, C)/(S, 0))$ gives us an extremal neighborhood in the sence of [14] (not necessary flipping). As in the introduction we apply the Minimal Model Program to (X^b, C^b) over $(S^b, 0)$:

$$\begin{array}{ccc} (X^b, C^b) & \xrightarrow{-p} & (Y, L) \\ & \searrow & \swarrow \\ & (S^b, 0) & \end{array}$$

Here $(Y, L) \rightarrow (S^b, 0)$ is a Mori conic bundle with $\rho(Y, S^b) = 1$. Then the map p contracts at least $d - 1$ divisors. Let E be a proper transforms one of them on X^b . Since f^b is flat $\Gamma := f^b(E)$ is a curve on S^b . Moreover for a general point $s \in \Gamma$ the preimage $f^{b^{-1}}(s)$ is a reducible conic, so $f^{b^{-1}}(s) = \ell_1 + \ell_2$. Therefore $f^{b^{-1}}(\Gamma)$ has exactly two irreducible components $E \supset \ell_1$ and $E' \supset \ell_2$. Consider the orbit $\{E = E_1, E_2, \dots, E_k\}$ of E under the action of $\mathbb{Z}_d$. Obviously, every $f^b(E_i)$ is a curve on S^b . We have $\sum E_i \sim aK_{X^b}$ for some $a \in \mathbb{Q}$ because $\mathbb{Z}_d$ -invariant part of $\text{Pic}(X^b) \otimes \mathbb{Q}$ is $\text{Pic}(X) \otimes \mathbb{Q} \simeq \mathbb{Q}$. But for a general fiber $\ell := f^{b^{-1}}(s)$, over $s \in S^b$ we have $(\ell \cdot \sum E_i) = 0$, $(\ell \cdot K_{X^b}) < 0$, so $a = 0$, and $\sum E_i \sim 0$. Further $(\ell_1 \cdot E) < 0$, $(\ell_1 \cdot E') > 0$. It gives us $(\ell_1 \cdot E_i) > 0$ for some $E_i \in \{E = E_1, E_2, \dots, E_k\}$. Then $E' = E_i$, i. e. there exists $\sigma \in \mathbb{Z}_d$ such that $\sigma(E) = E'$. From the symmetry we get that the orbit $\{E = E_1, E_2, \dots, E_k\}$ may be divided into couples of divisors E_j, E'_j such that $f^b(E_j) = f^b(E'_j)$ is a curve. Thus both k and d are even. $\square$

Lemma 3.7. *Let $f: (X, C \simeq \mathbb{P}^1) \rightarrow (S, 0)$ be a Mori conic bundle with irreducible central fiber and with a unique point P of index $m > 1$. Assume that P is an imprimitive of subindex $\bar{m}$ and splitting degree d (so $m = \bar{m}d$). Then*

- (i) $2\bar{m} \equiv 0 \pmod{d}$,
- (ii) $(-K_X \cdot C) = 1/\bar{m}$.

Corollary 3.7.1. *In conditions 3.7, $\bar{m} \geq d/2$. In particular, if $d \geq 3$, then $\bar{m} > 1$.*

Proof. Again consider the base change 3.1. Terminal singularities are rational and therefore Cohen-Macaulay [9]. By [12, Theorem 23.1], f^b is flat. Let X_s^b be a general fiber of f^b . Then $X_s^b \equiv nC^b$ for some $n \in \mathbb{N}$. We have

$$2 = (-K_{X^b} \cdot X_s^b) = n(-K_{X^b} \cdot C^b) = n(-K_{X^b} \cdot \sum C^b(i)) = nd(-K_{X^b} \cdot C^b(1)).$$

It is clear that $(-K_{X^\flat} \cdot C^\flat(1)) \in \frac{1}{\bar{m}}\mathbb{Z}$ because $-\bar{m}K_{X^\flat}$ is Cartier. If $(-K_{X^\flat} \cdot C^\flat(1)) = \delta/\bar{m}$ for some $\delta \in \mathbb{N}$, then $2 = nd\delta/\bar{m}$ or equivalently $2\bar{m} = nd\delta$. This proves (i). Since every $(X^\flat, C(i))$ is a primitive extremal neighborhood and $-K_{X^\flat}$ is a generator of $\text{Cl}^{\text{sc}}(X^\flat, P^\flat) \simeq \mathbb{Z}_{\bar{m}}$, $(\delta, \bar{m}) = 1$. Thus $\delta = 1$ by Lemma 3.6. On the other hand, $(-K_X \cdot C) = (-K_{X^\flat} \cdot C^\flat(1)) = \delta/\bar{m}$, because $K_{X^\flat} = g^*(-K_X)$. The rest is obvious. $\square$

Using Lemma 2.10 we obtain.

Corollary 3.7.2. *Let $f: (X, C \simeq \mathbb{P}^1) \rightarrow (S, 0)$ be a Mori conic bundle. Assume that $(S, 0)$ is singular and X contains the only one non-Gorenstein point P . Then in notations of 2.8.1 one has $\text{ord}(x_3) \equiv 1 \pmod{\bar{m}}$.*

$\square$

The following construction explains why Conjecture $(*_2)$ can be useful for studying Mori conic bundles.

3.8. Construction (double cover trick [7]). Let $f: (X, C) \rightarrow (S, 0)$ be a Mori conic bundle and let $D \in |-2K_X|$ be a general member. Assume that $K_X + 1/2D$ is log-terminal (i. e. $(*_2)$ holds). Then there exists a double cover $h: Y \rightarrow X$ with ramification divisor D such that Y has only canonical Gorenstein singularities (see [7, 8.5]). Thus we have two morphisms

$$Y \xrightarrow{h} X \xrightarrow{f} S.$$

By the Hurwitz formula, $K_Y = h^*(K_X + 1/2D) = 0$. Therefore the composition map $g: Y \rightarrow S$ gives us an elliptic fiber space structure on Y .

4. THE MAIN RESULT

The main result of this section is the following

Theorem 4.1. *Let $f: (X, C \simeq \mathbb{P}^1) \rightarrow (S, 0)$ be a Mori conic bundle with only one non-Gorenstein point P . Then one of the following holds*

4.1.1. *$(S, 0)$ is non-singular.*

4.1.2. *$(S, 0)$ is a DuVal point of type A_1 and splitting degree of P is 2 and X contains at most one more singular (Gorenstein) point. There are the following subcases:*

- (i) (cf. [18], [19]) *$f: (X, C) \rightarrow (S, 0)$ is a quotient of a conic bundle $f^\flat: (X^\flat, C^\flat) \rightarrow (\mathbb{C}^2, 0)$ by $\mathbb{Z}_2$, P is the only singular point. In some coordinate system $(x, y, z; u, v)$ in $\mathbb{P}^2 \times \mathbb{C}^2$ the variety $X^\flat$ can be defined by the equation*

$$x^2 + y^2 + \psi(u, v)z^2 = 0, \quad \psi(u, v) \in \mathbb{C}\{u, v\},$$

where $\psi(u, v)$ has no multiple factors and consists only of monomials of even degree. Here $X^\flat \rightarrow \mathbb{C}^2$ is the natural projection and the action of $\mathbb{Z}_2$ has the form

$$u \rightarrow -u, \quad v \rightarrow -v, \quad x \rightarrow -x, \quad y \rightarrow y, \quad z \rightarrow z.$$

Moreover

- (a) if $\text{mult}_{(0,0)} \psi(u, v) = 2$, then (X, P) is of type $cA/2$,
 - (b) if $\text{mult}_{(0,0)} \psi(u, v) \geq 4$, then (X, P) is of type $cAx/2$.
- (ii) (X, P) is of type $cAx/4$, more precisely

$$(X, P) \simeq (\{y_1^2 - y_2^2 + y_3\varphi_3 + y_4\varphi_4 = 0\}/\mathbb{Z}_4(1, 3, 3, 2),$$

$$C = \{y_1^2 - y_2^2 = y_3 = y_4 = 0\}/\mathbb{Z}_4,$$

where φ_3, φ_4 are semi-invariants with weights

$$\text{wt}(\varphi_3) \equiv 3 \pmod{4}, \quad \text{wt}(\varphi_4) \equiv 0 \pmod{4}$$

(this is a point of type $II^\vee$ in Mori's classification [14]).

- (iii) (X, P) is the cyclic quotient singularity $\mathbb{C}^3/\mathbb{Z}_4(1, 3, 1)$ and

$$C = \{y_1^2 - y_2^2 = y_3 = 0\}/\mathbb{Z}_4.$$

- (iv) $\bar{m}$ is even and ≥ 4 , (X, P) is the cyclic quotient singularity $\mathbb{C}^3/\mathbb{Z}_{2\bar{m}}(1, -1, \bar{m} + 1)$

$$C = \{y_2^2 - y_1^{2\bar{m}-2} = y_3 = 0\} \ni 0)/\mathbb{Z}_{2\bar{m}}$$

(this is a point of type $IC^\vee$ in [14], see also example 2.4).

- (v) $\bar{m}$ is even, (X, P) is of type cA/m , where $m = 2\bar{m}$. More precisely

$$(X, P) \simeq \{y_1y_2 + y_2\varphi_1 + y_3\varphi_2 + (y_4^2 - y_1^{2\bar{m}})\varphi_3 = 0\}/\mathbb{Z}_{2\bar{m}}(1, -1, \bar{m} + 1, 0)$$

and

$$C = \{y_2 = y_3 = y_4^2 - y_1^{2\bar{m}} = 0\}/\mathbb{Z}_{2\bar{m}},$$

where $\varphi_i, i = 1, 2, 3$ are semi-invariants with weights

$$\text{wt}(\varphi_1, \varphi_2, \varphi_3) \equiv (1, \bar{m} - 1, 0) \pmod{2\bar{m}}.$$

4.1.3. $(S, 0)$ is a DuVal point of type A_3 , (X, P) is a cyclic quotient singularity of type $\mathbb{C}^3/\mathbb{Z}_8(a, -a, 1)$, splitting degree of P is 4 and X has no other singular points. There are two subcases

- (i) $a = 1$, $C = \{y_1^4 - y_2^4 = y_3 = 0\}/\mathbb{Z}_8$,
- (ii) $a = 3$, $C = \{y_2^2 - y_3^2 = y_2y_3 - y_1^2 = 0\}/\mathbb{Z}_8$.

Remark 4.2. We will show that in 4.1.2, 4.1.3 a general member of $|-K_X|$ does not contain C except 4.1.2 (iv), (v). In these two cases Conjecture $(*_2)$ holds. Using the same arguments one can prove that in 4.1.1 at least one of Conjectures $(*_1)$, $(*_2)$ or $(*_3)$ holds. More precise results for the case of non-singular base surface will be published elsewhere.

Before starting the proof we consider examples of Mori conic bundles such as in 4.1.2 (ii), (iii) and 4.1.3. Unfortunately I don't know examples of Mori conic bundles as in 4.1.2 (iv) or (v).

Example 4.3. Let $Y \subset \mathbb{P}_{x,y,z,t}^3 \times \mathbb{C}_{u,v}^2$ be a non-singular subvariety given by the equations

$$\begin{cases} xy = ut^2 \\ (x + y + z)z = vt^2 \end{cases}$$

It is easy to check that Y is non-singular. The projection $g: Y \rightarrow \mathbb{C}^2$ gives us an elliptic fibration whose fibers are intersections of two quadrics in $\mathbb{P}^3$. The special fiber $g^{-1}(0)$ is the union of four lines meeting at one point $Q := \{(x, y, z, t; u, v) = (0, 0, 0, 1; 0, 0)\}$. Consider also the following action of $\mathbb{Z}_8$ on Y :

$$\begin{aligned} x &\rightarrow \varepsilon^{-3}z, & y &\rightarrow \varepsilon(x + y + z), & z &\rightarrow -\varepsilon y, & t &\rightarrow t, \\ & & & & u &\rightarrow \varepsilon^{-2}v, & v &\rightarrow -\varepsilon^2u, \end{aligned}$$

where $\varepsilon := \exp(2\pi i/8)$. Then a generator of $\mathbb{Z}_8$ has only one fixed point Q and the quotient (Y, Q) is terminal of type $\frac{1}{8}(1, -1, 3)$. The locus of fixed point for $\mathbb{Z}_2 \subset \mathbb{Z}_8$ consists of the point Q and the divisor $B := \{t = 0\} \cap Y$. Let $h: Y \rightarrow Y' := Y/\mathbb{Z}_2$ be the quotient morphism. By the Hurwitz formula $0 = K_Y = h^*K_{Y'} + B$, hence $h^*(-K_{Y'}) = B$. Therefore $-K_{Y'}$ is ample over $\mathbb{C}^2$ and $Y' \rightarrow \mathbb{C}^2$ is a Mori conic bundle with singular point $h(Q)$ of type $\frac{1}{2}(1, 1, 1)$ and reducible central fiber. Note that

$$g: Y \xrightarrow{h} X \xrightarrow{f} S.$$

is the double cover trick 3.8. Further the induced action $\mathbb{Z}_4 = \mathbb{Z}_8/\mathbb{Z}_2$ on Y' is free outside point $h(Q)$ and permutes components of the central fiber. Therefore the anticanonical divisor of $X := Y'/\mathbb{Z}_4$ is ample over $\mathbb{C}^2/\mathbb{Z}_4$ and the central fiber of the projection $X \rightarrow \mathbb{C}^2/\mathbb{Z}_4$ is irreducible. Finally we obtain the Mori conic bundle $X := Y/\mathbb{Z}_8 \rightarrow \mathbb{C}^2/\mathbb{Z}_4$ such as in 4.1.3. The diagram

$$\begin{array}{ccc} Y' & \longrightarrow & X \\ f' \downarrow & & \downarrow f \\ \mathbb{C}^2 & \longrightarrow & \mathbb{C}^2/\mathbb{Z}_4 \end{array}$$

is exactly as in the construction 3.1.

Example 4.4. Let $Y \subset \mathbb{P}_{x,y,z,t}^3 \times \mathbb{C}_{u,v}^2$ is given by the equations

$$\begin{cases} xy = ut^2 \\ z^2 = u(x^2 + y^2) + vt^2 \end{cases}$$

Consider the action of $\mathbb{Z}_4$ on Y :

$$x \rightarrow y, \quad y \rightarrow -x, \quad z \rightarrow iz, \quad t \rightarrow t, \quad u \rightarrow -u, \quad v \rightarrow -v.$$

As above $X := Y/\mathbb{Z}_4 \rightarrow \mathbb{C}^2/\mathbb{Z}_2$ is a Mori conic bundle such as in 4.1.2 (iii) with unique singular point of type $\frac{1}{4}(1, 1, 3)$

Example 4.5. Let $Y \subset \mathbb{P}_{x,y,z,t}^3 \times \mathbb{C}_{u,v}^2$ is given by the equations

$$\begin{cases} xy = (u^{2k+1} + v)t^2, \\ z^2 = u(x^2 - y^2) + vt^2 \end{cases} \quad k \in \mathbb{N}$$

Define the action of $\mathbb{Z}_4$ on Y by

$$x \rightarrow iy, \quad y \rightarrow ix, \quad z \rightarrow iz, \quad t \rightarrow t, \quad u \rightarrow -u, \quad v \rightarrow -v.$$

Then $X := Y/\mathbb{Z}_4 \rightarrow \mathbb{C}^2/\mathbb{Z}_2$ is a Mori conic bundle such as in 4.1.2 (ii) with unique singular point of type $cAx/4$.

4.6. To begin the proof of Theorem 4.1 we suppose that $(S, 0)$ is singular (otherwise we have case 4.1.1). Then by 3.5 the point P is imprimitive. Denote by m , d and $\bar{m}$ its index, splitting degree and subindex, respectively. First we prove the existence of good divisors in $|-K_X|$ (with two exceptions 4.1.2 (iv), (v)). As in [14, §7] we will use local methods to find good divisors in $|-K_X|$ or $|-2K_X|$.

Lemma 4.7. (cf. [14, proof of 7.3]). *Let $f: (X, C \simeq \mathbb{P}^1) \rightarrow (S, 0)$ be a Mori conic bundle with only one non-Gorenstein point P . Let $\pi: (X^\sharp, P^\sharp) \rightarrow (X, P)$ be the canonical cover and $X^\sharp \subset \mathbb{C}^4$ be an embedding as in 1.2.1. Then a general member $F \in |-K_X|$ does not contain C and has only DuVal singularity at P iff we have $\text{ord}(x_3) < \bar{m}$ in a normalized coordinate system $(x_1, \dots, x_4)$ in $\mathbb{C}^4$. (By 3.7.2, $\text{ord}(x_3) < \bar{m}$ implies $\text{ord}(x_3) = 1$, if $(S, 0)$ is singular.)*

Proof. Assume that $\text{ord}(x_3) < \bar{m}$. From Lemma 3.7 we have $(-K_X \cdot C) = 1/\bar{m}$. Take a general member $F \in |-K_{(X,P)}|$. Proposition 2.10 yields $(F \cdot C)_P = \text{ord}(x_3)/\bar{m} < 1$. Then $F + K_X$ can be considered as a Cartier divisor on X and in our case $-1 < ((F + K_X) \cdot C) < 1$. Therefore $(F \cdot C) = (-K_X \cdot C)$. Since $\text{Pic}(X) \simeq \mathbb{Z}$, $F \in |-K_X|$ and by Theorem 1.2.2, F has only DuVal singularity at P . The inverse implication is obvious. $\square$

Thus it is sufficient to show only $\text{ord}(x_3) < \bar{m}$. First we consider the exceptional series of terminal points 1.2.1, (ii).

Lemma 4.8. (cf. [14, 4.2]). *If $(X, P) \supset C$ is an imprimitive point of type $cAx/4$, then $\bar{m} = 2$ and*

	x_1	x_2	x_3	x_4	
wt	1	3	3	2	mod 4
ord	1	1	1	2	

Proof. (cf. [14, 3.8]) From Corollary 3.7.1, one has $\bar{m} = 2$. By 2.8.1, $\text{ord}(x_4) \equiv 0 \pmod{\bar{m}}$ and there exists an invariant monomial ψ such that $\text{ord}(\psi) = \bar{m} = 2$. Since $\text{wt}(x_i) \not\equiv 0 \pmod{4}$ for $i = 1, 2, 3, 4$, we have $\psi = x_i x_j$. Up to permutation $\{1, 2\}$, we may assume that $\text{wt}(x_2) \equiv \text{wt}(x_3)$. Therefore $\text{ord}(x_2) = \text{ord}(x_3)$. It gives us $\psi = x_1 x_2$ or $x_1 x_3$, whence $\text{ord}(x_1) = \text{ord}(x_2) = \text{ord}(x_3) = 1$. If necessary we can replace the character χ with $-\chi$ to obtain $\text{wt}(x_1) \equiv 1$. Finally $\text{wt}(x_4) \equiv \text{wt}(x_1^2) \equiv 2 \pmod{m}$ gives as $\text{ord}(x_4) = 2$. $\square$

Since $\text{ord}(x_3) = 1$, we have a good divisor in $|-K_X|$ in this case. By 2.8.1, components $C^\sharp(1), C^\sharp(2)$ of curve $C^\sharp$ are images of

$$t \longrightarrow (t, t, t, t^2), \quad t \longrightarrow (it, -it, -it, -t^2).$$

Therefore $C^\sharp$ can be given by the equations $x_4 - x_1^2 = x_2 - x_3 = x_2^2 - x_1^2 = 0$. By changing coordinates as $y_1 = x_1, y_2 = x_2, y_3 = x_3 - x_1^2, y_4 = x_4 - x_1^2$ we obtain case 4.1.2 (ii) of our theorem.

From now we assume that (X, P) is from the main series. In particular $\text{wt}(x_4) \equiv 0 \pmod m, \text{ord}(x_4) = \bar{m}$.

Lemma 4.9. *Let $f(X, C \simeq \mathbb{P}^1) \rightarrow (S, 0)$ be a Mori conic bundle. Assume that $P \in X$ is the only point of index $m > 1$. Assume further that (X, C) is imprimitive at P of splitting degree d . Then one of the following holds*

4.9.1. *A general member of $|-K_X|$ does not contain C and has DuVal singularity at P . (By 4.7, this is equivalent to $\text{ord}(x_3) = 1$).*

4.9.2. *$d = 2, \bar{m}$ is even a general member of $|-2K_X|$ does not contain C , and up to permutation $x_1, \dots, x_4$ in some normalized coordinate system we have*

$$\begin{array}{llllll} \text{(i)} & & x_1 & x_2 & x_3 & x_4 & \\ & \text{ord} & 1 & \bar{m} - 1 & \bar{m} + 1 & \bar{m} & \\ & \text{wt} & 1 & -1 & \bar{m} + 1 & 0 & \pmod m \end{array} \quad \bar{m} \geq 4 \quad \text{or}$$

$$\begin{array}{llllll} \text{(ii)} & & x_1 & x_2 & x_3 & x_4 & \\ & \text{ord} & 1 & 2\bar{m} - 1 & \bar{m} + 1 & \bar{m} & \\ & \text{wt} & 1 & -1 & \bar{m} + 1 & 0 & \pmod m \end{array}$$

4.10. **Proof.** Denote $a_i := \text{ord}(x_i)$. Since $(\text{wt}(x_i), m) = 1$, for $i = 1, 2, 3$, one has $(a_i, \bar{m}) = 1$, $i = 1, 2, 3$. For 4.9.2 it is sufficient to prove that $a_3 < \bar{m}$. Let $a_3 > \bar{m}$. First suppose that $\text{ord}(x_1^m) = 2\bar{m}$. We claim that in this situation 4.9.2 holds. Indeed, then $a_1 m = a_1 \bar{m} d = 2\bar{m}$, hence $d = 2, a_1 = 1, m = 2\bar{m}$. Thus $\text{wt}(x_1) \equiv \text{wt}(x_3) \equiv 1 \pmod{\bar{m}}$. By the assumptions in 2.8.1 and because $a_3 > \bar{m}$, $\text{wt}(x_1) \not\equiv \text{wt}(x_3) \pmod m$. Since $(\text{wt}(x_1), m) = (\text{wt}(x_3), m) = 1$, $\bar{m}$ is even. Changing a generator of $\mathbb{Z}_m$ if necessary we may achieve that $\text{wt}(x_1) \equiv 1 \pmod m$. The rest follows from normalizedness. Taking into account Lemma 3.6, from now we may assume that $\text{ord}(x_1^m) > 3\bar{m}$, and by symmetry, $\text{ord}(x_2^m) > 3\bar{m}$. We will derive a contradiction with $a_3 > \bar{m}$.

4.11. By Corollaries 2.6.1 and 2.11.4, there exist different simple invariant monomials ψ_1, ψ_2, ψ_3 such that

$$3\bar{m} \geq i_P \bar{m} \geq \text{ord}(\psi_1) - a_1 + \text{ord}(\psi_2) - a_2 + \text{ord}(\psi_3) - a_3 - \bar{m} w_P.$$

Remind that the last inequality can be rewritten as

$$3\bar{m} \geq i_P \bar{m} \geq \text{ord}(\nu_1) + \text{ord}(\nu_2) + \text{ord}(\nu_3) - \bar{m} w_P,$$

where $\nu_i x_i = \psi_i$, $i = 1, 2, 3$. In particular $\text{ord}(\nu_1) + \text{ord}(\nu_2) \leq 3\bar{m}$. Since $\text{wt}(\nu_1) \equiv \text{wt}(x_2) \pmod m, \text{wt}(\nu_2) \equiv \text{wt}(x_1) \pmod m$, by 2.8.1, we have $a_1 + a_2 \leq 3\bar{m}$. Consider the following cases.

4.12. **Case** $a_1 + a_2 = 3\bar{m}$. Then we have $\text{ord}(\nu_1) + \text{ord}(\nu_2) = 3\bar{m}$. Therefore $i_P = 3$, $w_P < 1$ (by the inequality in 2.6.1) and $\text{ord}(\nu_3) = \bar{m}w_P < \bar{m}$ by 4.11. Since $a_3 > \bar{m}$, the monomial ν_3 does not depend on x_3 . Up to permutation of x_1, x_2 we may assume that $\nu_3 = x_1^\alpha$ for some $\alpha \in \mathbb{N}$ such that $\alpha \text{wt}(x_1) + \text{wt}(x_3) \equiv 0 \pmod{m}$ and $\alpha a_1 < \bar{m}$. Thus $\psi_3 = x_1^\alpha x_3$ and $a_1 < \bar{m}$, $a_2 > 2\bar{m}$.

Further from normalizedness and from $\text{wt } \nu_1 \equiv \text{wt } x_2 \pmod{m}$, we have $\text{ord}(\nu_1) = a_2 > 2\bar{m}$, so $\text{ord}(\nu_2) = a_1 < \bar{m}$. It gives us $\nu_2 = x_1$, $\psi_2 = x_1 x_2$ and $\text{ord}(\psi_1) = 3\bar{m}$. By the assumption in 4.10, ψ_1 is not a power of x_1 . If ν_1 depends on x_2 , then $\psi_1 = x_1 x_2 = \psi_2$, a contradiction. Hence $\psi_1 = x_1^\beta x_3^\gamma$, where $\beta, \gamma > 0$, $a_1\beta + a_3\gamma = 3\bar{m}$ and $\text{wt}(x_1)\beta + \text{wt}(x_3)\gamma \equiv 0 \pmod{m}$. Since $a_3 > \bar{m}$, $\gamma \leq 2$. On the other hand if $\gamma = 1$, then $x_1^{\beta-\alpha} = \psi_1/\psi_3$ is an invariant of order $\leq 2\bar{m}$, a contradiction with assumptions in 4.10. Thus $\gamma = 2$. It gives us $a_1\beta + 2a_3 = 3\bar{m}$ and $a_1\alpha + a_3 = 2\bar{m}$. Then $(2\alpha - \beta)a_1 = \bar{m}$, so $a_1 = 1$ and $2\alpha - \beta = \bar{m}$, because $(a_1, \bar{m}) = 1$. On the other hand $\beta \text{wt}(x_1) + 2\text{wt}(x_3) \equiv 0 \pmod{m}$ and $\alpha \text{wt}(x_1) + \text{wt}(x_3) \equiv 0 \pmod{m}$, so $\bar{m} \text{wt}(x_1) = (2\alpha - \beta) \text{wt}(x_1) \equiv 0 \pmod{m}$, the contradiction with $(m, \text{wt}(x_1)) = 1$.

4.13. Thus $a_1 + a_2 \leq 2\bar{m}$. Then there are no invariant monomials in x_1, x_2, x_3 of order $< \text{ord}(x_1 x_2)$. Up to permutation of ψ_1 and ψ_2 we can take $\psi_1 = x_1 x_2$. Since ψ_2 is simple, it depends only on x_2, x_3 .

4.14. **Case** $a_1 + a_2 = 2\bar{m}$. Simple invariant monomials of order $\leq 2\bar{m}$ may be only of the form $x_1 x_2$ or $x_i^\alpha x_3$, $i = 1, 2$ (if $\alpha \text{wt}(x_i) + \text{wt}(x_3) \equiv 0 \pmod{m}$ and $\alpha a_i + a_3 = 2\bar{m}$). On the other hand by the inequality in 4.11,

$$\text{ord}(\psi_2) \leq 3\bar{m} + a_1 + a_2 - \text{ord}(\psi_1) = 3\bar{m}.$$

There are two subcases.

4.14.1. *Subcase* $\text{ord}(\psi_2) = 2\bar{m}$. Then $\psi_2 = x_2^\alpha x_3$, because ψ_2 depends on x_2 . Therefore $a_3 < 2\bar{m}$, $a_2\alpha < \bar{m}$ and $a_1 > \bar{m}$. Further, if we take in 2.9 $\psi = x_2^\alpha$, we obtain $\bar{m}w_P = \text{ord}(x_2^\alpha) = a_2\alpha < \bar{m}$. It follows from inequality 4.11 that $\text{ord}(\psi_3) \leq \bar{m}w_P + a_3 + \bar{m} \leq 3\bar{m}$, hence $\text{ord}(\psi_3) = 3\bar{m}$. Using $\partial\psi_3/\partial x_3 \neq 0$ we get $\psi_3 \in \{x_1 x_3, x_2^\gamma x_3, x_2^\gamma x_3^2\}$. But if $\psi_3 = x_1 x_3$, then $\text{wt}(x_3) \equiv \text{wt}(x_2) \pmod{m}$, and by normalizedness $a_3 = a_2 < \bar{m}$, a contradiction.

If $\psi_3 = x_2^\gamma x_3$, then $\text{ord}(x_2^{\gamma-\alpha}) = \text{ord}(\psi_3/\psi_2) = \bar{m}$. We derive a contradiction with assumptions in 4.10.

Thus $\psi_3 = x_2^\gamma x_3^2$, where $\gamma \text{wt}(x_2) + 2\text{wt}(x_3) \equiv 0 \pmod{m}$ and $\gamma a_2 + 2a_3 = 3\bar{m}$. Consider the monomial $\nu = x_2^{\alpha-\gamma} = x_2 \psi_2 / \psi_3$. We have $\text{ord}(\nu) = a_3 - \bar{m} < a_3$. On the other hand $\text{wt}(\nu) \equiv \text{wt}(x_3) \pmod{m}$. Hence $\text{wt}(\nu) \equiv \text{wt}(x_3) \pmod{m}$ and $\text{ord}(\nu) < \text{ord}(x_3)$, a contradiction with normalizedness.

4.14.2. *Subcase* $\text{ord}(\psi_2) = 3\bar{m}$. By the inequality in 4.11, $i_P = 3$, $\text{ord}(\psi_3) = a_3 + \bar{m}w_P$ and by Corollary 2.6.1, $w_P < 1$. Therefore $\text{ord}(\psi_3) < a_3 + \bar{m}$. Thus one has $\psi_2 \in \{x_2^\beta x_3^2, x_2^\beta x_3\}$. But if $\psi_2 = x_2^\beta x_3^2$, then $a_2\beta + 2a_3 = 3\bar{m}$. Hence $a_3 < 2\bar{m}$, $a_2 < \bar{m}$, $a_1 > \bar{m}$ and $\text{ord}(\psi_3) = 2\bar{m}$. So $\text{ord}(\psi_3) = 2\bar{m}$

and ψ_3 depends on x_2, x_3 . Then we can permute ψ_2, ψ_3 and get the subcase above.

Therefore $\psi_2 = x_2^\beta x_3$. Then $a_2\beta + a_3 = 3\bar{m}$, $a_3 < 3\bar{m}$. We claim that $a_2 < \bar{m}$, $a_1 > \bar{m}$. Indeed if otherwise, we have $\beta = 1$, $\text{wt}(x_3) \equiv \text{wt}(x_1)$ and by normalizedness $a_3 = a_1 < \bar{m}$, a contradiction. Further $\text{ord}(\psi_3) \leq 3\bar{m}$ gives us

$$\psi_3 \in \{x_2^\gamma x_3, \quad x_2^\gamma x_3^2, \quad x_1 x_3\}.$$

If $\psi_3 = x_2^\gamma x_3$, then the monomial $\psi_2/\psi_3 = x_2^{\beta-\gamma}$ is invariant and has order $\leq \bar{m}$, a contradiction with assumptions in 4.10. But if $\psi_3 = x_2^\gamma x_3^2$, then we can permute ψ_2, ψ_3 and derive a contradiction as above. Therefore ψ_3 depends on x_1, x_3 , so $\psi_3 = x_1 x_3$. Again from normalizedness we have $a_3 = a_2 < \bar{m}$, a contradiction.

4.15. Case $a_1 + a_2 = \bar{m}$. Recall that $\psi_1 = x_1 x_2$. As above by the inequality in 4.11

$$\text{ord}(\psi_2) \leq 3\bar{m},$$

and if the equality holds, then $i_P = 3$, $w_P < 1$ and

$$\text{ord}(\psi_3) = a_3 + \bar{m}w_P < a_3 + \bar{m}.$$

Since $\psi_1 = x_1 x_2$ and ψ_2 are different simple invariant monomials, we have that ψ_2 depends on x_3 . Then obviously $\text{ord}(\psi_2) \geq 2\bar{m}$. Consider subcases.

4.15.1. Subcase $\text{ord}(\psi_2) = 2\bar{m}$. Then it is easy to see $\psi_2 = x_2^\beta x_3$. It implies that

$$a_2\beta + a_3 = 2\bar{m}, \quad \text{wt}(x_1)\beta \equiv \text{wt}(x_3) \pmod{m}.$$

Moreover we may assume that $\beta \geq 2$ (otherwise we derive a contradiction as in 4.14.1). If we take in 2.9 $\psi = \psi_2/x_3 = x_2^\beta$, we get $w_P < 1$. Thus $\text{ord}(\psi_3) \leq \bar{m} + \bar{m}w_P + a_3 < 3\bar{m} + \bar{m}w_P$ and $\text{ord}(\psi_3) \leq 3\bar{m}$. We claim that $\text{ord}(\psi_3) = 3\bar{m}$. Indeed, if otherwise, we have $\psi_3 = x_1^\gamma x_3$, where $a_1\gamma + a_3 = 2\bar{m}$. It gives us $a_1(\gamma - 1) + a_2(\beta - 1) + 2a_3 = 3\bar{m}$. Since $\beta \geq 2$, $\gamma = 1$. But then $x_1 x_3$ is an invariant and $a_3 = a_2 < \bar{m}$, a contradiction.

Thus $\text{ord}(\psi_3) = 3\bar{m}$. Taking into account that ψ_3 and ψ_3/ψ_2 are not powers of x_i , $i = 1, \dots, 4$, we obtain

$$\psi_3 \in \{x_1^\gamma x_3, \quad x_1^\gamma x_3^2, \quad x_2^\gamma x_3^2\}.$$

If $\psi_3 = x_1^\gamma x_3$, then $a_1\gamma + a_3 = 3\bar{m}$, so $a_3 = 3\bar{m} - a_1\gamma$ and $a_1(\gamma + \beta) = \bar{m}(\beta + 1)$. Since $(a_1, \bar{m}) = 1$, $\gamma + \beta = \bar{m}k$, where k is a positive integer. Then $\beta = a_1\bar{m}k - 1 \geq \bar{m} - 1$. The equality $a_2\beta + a_3 = 2\bar{m}$ is possible only if $a_1 = a_2 = 1$, $\bar{m} = 2$, but then $\beta = 1$, a contradiction.

Assume that $\psi_3 = x_1^\gamma x_3^2$. Then $a_1\gamma + 2a_3 = 3\bar{m}$, $a_1(\gamma - 1) + a_2(\beta - 1) + 3a_3 = 4\bar{m}$. Since $\beta \geq 2$, $\gamma = 1$. Whence

$$3\bar{m} = a_1 + 2a_3 = a_1 + 2(2\bar{m} - a_2\beta) = 4\bar{m} + a_1 - 2a_2\beta = (4 - 2\beta)\bar{m} + (1 + 2\beta)a_1.$$

Thus $a_1(2\beta + 1) = \bar{m}(2\beta - 1)$. Since $(a_1, \bar{m}) = 1$, it implies $\bar{m} = 2\beta + 1$, $a_1 = 2\beta - 1$. On the other hand $\text{wt}(x_1) + 2\text{wt}(x_3) \equiv 0 \pmod{m}$ and $\text{wt}(x_1)\beta \equiv$

$\text{wt}(x_3) \bmod m$ imply $\bar{m} = 1 + 2\beta \equiv 0 \bmod m$. So $\text{wt}(x_1)(1 + 2\beta) \equiv 0 \bmod m$, a contradiction.

Finally assume that $\psi_3 = x_2^\gamma x_3^2$. Then $a_2\gamma + 2a_3 = 3\bar{m}$ and $a_2\beta + a_3 = 2\bar{m}$. It gives us $\bar{m} = a_2(2\beta - \gamma)$, so $a_2 = 1$, $\bar{m} = 2\beta - \gamma$. On the other hand, $\text{wt}(x_1)\gamma \equiv 2\text{wt}(x_2) \bmod m$, because ψ_3 is an invariant. Whence $2\text{wt}(x_1)\beta \equiv \text{wt}(x_1)\gamma \bmod m$ and $\bar{m} = 2\beta - \gamma \equiv 0 \bmod m$, a contradiction.

4.15.2. *Subcase* $\text{ord}(\psi_2) = 3\bar{m}$. Since ψ_2 depends on x_2 and x_3 , $a_3 < 3\bar{m}$. We claim that $a_3 < 2\bar{m}$. Indeed, if otherwise, we have $\psi_2 = x_2^\beta x_3$ and $\text{ord}(\psi_3) = 3\bar{m}$, hence $\psi_3 = x_1^\alpha x_3$. Thus $a_1\alpha + a_3 = 3\bar{m}$ and $a_2\beta + a_3 = 3\bar{m}$ give us $(\alpha - 1)a_1 + (\beta - 1)a_2 + 2a_3 = 5\bar{m}$. So $\alpha = 1$ or $\beta = 1$. But then x_1x_3 or x_2x_3 is an invariant. By normalizedness $a_3 < \bar{m}$, a contradiction.

Therefore $a_3 < 2\bar{m}$, hence $\text{ord}(\psi_3) = 2\bar{m}$,

$$\psi_3 \in \{x_1^\gamma x_3, \quad x_2^\gamma x_3\}, \quad \psi_2 \in \{x_2^\beta x_3, \quad x_2^\beta x_3^2\}.$$

But then we can permute $\{\psi_2, \psi_3\}$ and $\{x_1, x_2\}$ and get the previous subcase. This proves Lemma 4.9. $\square$

4.16. **Proof of Theorem 4.1.** First we treat 4.9.2 (cf. [14, (4.4)]). We claim that in case (i) of 4.9.2 (X, P) is a cyclic quotient singularity. Let $C^\sharp = C^\sharp(1) \cup C^\sharp(2)$ be irreducible decomposition of $C^\sharp = (\pi^{-1}(C))_{\text{red}}$. The component $C^\sharp(1)$ is the image of $t \longrightarrow (t, t^{\bar{m}-1}, t^{\bar{m}+1}, t^{\bar{m}})$. The curve $C^\sharp(1)$ is a complete intersection:

$$C^\sharp(1) = \{x_1x_2 - x_4 = x_3 - x_1^{\bar{m}+1} = x_2 - x_1^{\bar{m}-1} = 0\}$$

and so is $C^\sharp$:

$$C^\sharp = \{x_1x_2 - x_4 = x_3 - x_1^{\bar{m}+1} = x_2^2 - x_1^{2\bar{m}-2} = 0\}.$$

Therefore the equation of $X^\sharp$ can be written as

$$\phi = (x_1x_2 - x_4)\varphi_1 + (x_3 - x_1^{\bar{m}+1})\varphi_2 + (x_2^2 - x_1^{2\bar{m}-2})\varphi_3 = 0,$$

where $\varphi_i = \varphi_i(x_1, x_2, x_3, x_4)$ are semi-invariants. Since $\text{wt}(x_3) \not\equiv \pm \text{wt}(x_1), \text{wt}(x_2)$ and because (X, P) is of type cA/m , the equation of ϕ must contain either x_1x_2 or x_4 (see [13]). But this is possible only if $\varphi_1(0, 0, 0, 0) \neq 0$. Thus ϕ always contains x_4 and $X^\sharp$ is non-singular. We obtain case (iv) of 4.1.2. Case (ii) of 4.9.2 is treated by the similar way.

4.17. From now we will suppose that f is not from 4.1.2 (iv) or (v). Consider the base change from 3.1

$$\begin{array}{ccc} X^\flat & \xrightarrow{g} & X \\ f^\flat \downarrow & & \downarrow f \\ S^\flat & \xrightarrow{h} & S \end{array}$$

If $\bar{m} = 1$, then $X^\flat$ is Gorenstein and $f^\flat$ is a "classical" conic bundle [3] (it means that $S^\flat$ is non-singular and every fiber of $f^\flat$ is a conic in $\mathbb{P}^2$). Since

(X, C) is imprimitive, the central fiber C^b has exactly two components. The group $\mathbb{Z}_2$ permutes these components. In this case it is not difficult to write down an equation of X^b explicitly (see [18] or [19]). We obtain case 4.1.2, (i).

Assume now that $\bar{m} > 1$. By Lemma 4.9, the general member $F \in |-K_X|$ does not contain P and has DuVal point at P . The cover $f_F: (F, P) \rightarrow (S, 0)$ is finite of degree 2. Thus $(S, 0)$ is a quotient of (F, P) by an involution.

Proposition 4.17.1. (see [1]) *Let (F, P) be a germ of DuVal singularity and $\tau: (F, P) \rightarrow (F, P)$ be an (analytic) involution. Then up to analytic isomorphism there are only the following possibilities.*

no.	(F, P)	$(F, P)/\tau$	no.	(F, P)	$(F, P)/\tau$
1)	any	smooth	6)	A_{2k+1}	$\mathbb{C}^2/\mathbb{Z}_{4k+4}(1, 2k+1)$
2)	A_{2k+1}	D_{k+3}	7)	E_6	A_2
3)	E_6	E_7	8)	A_{2k}	$\mathbb{C}^2/\mathbb{Z}_{2k+1}(1, 2k-1)$
4)	D_k	D_{2k-2}	9)	D_k	A_1
5)	A_k	A_{2k+1}	10)	A_{2k+1}	A_k

Remark. In cases 6) and 8) the dual graphs of corresponding minimal resolutions of $(F, P)/\tau$ are

$$6) \quad \begin{array}{c} -4 \\ \circ \end{array} \quad (\text{for } k = 0),$$

$$6) \quad \underbrace{\begin{array}{ccccccc} -3 & & -2 & & -2 & & -3 \\ \circ & - & \circ & \dots & \circ & - & \circ \end{array}}_{k+1} \quad (\text{for } k > 0)$$

$$8) \quad \underbrace{\begin{array}{ccccccc} -2 & & -2 & & -2 & & -2 & & -3 \\ \circ & - & \circ & \dots & \circ & - & \circ & - & \circ \end{array}}_k$$

As noticed by Shokurov, these singularities are exactly all log-terminal singularities with discrepancies $\geq -1/2$.

4.17.2. *Proof of Theorem 4.1 (continue).* Since $(S, 0) = (F, P)/\tau$ is a cyclic quotient singularity by Proposition 3.2, cases 1)-4) are impossible. Further by assumptions in 4.17 we have $\bar{m} > 1$, hence $m > d$. Remind that in our situation d is the order of local fundamental group $\pi_1(S - \{0\})$ (see 3.5). In particular, $d = n + 1$ if $(S, 0)$ is a point of type A_n . Thus in case 5) we have $m > d = 2k + 2 \geq 4$. But (F, P) is a point of type A_k and has the topological index $k + 1$. This contradicts to 1.2.7. The same arguments show that the cases 6), 7), 8) are also impossible (if $\bar{m} > 1$). In case 9) $d = 2$. From the list

of 1.2.3 we see that $m > 2$ only if (X, P) is of type $cAx/4$ and then $\bar{m} = 2$, $m = 4$. This is case (ii) of 4.1.2 by Lemma 4.8.

Finally consider case 10). Then $d = k+1$. In the classification 1.2.3 (X, P) is of type cA/n . Point (F, P) has topological index $2k+2$ and $d = k+1$, so by Lemma 1.2.7 $(k+1)\bar{m} = d\bar{m} = m \leq 2(k+1)$, whence $\bar{m} = 2$. Moreover (X, P) is a cyclic quotient singularity in this case (because an equality in 1.2.7 holds). Lemma 3.7 gives us then $d = 2$ or 4 .

Suppose that $d = 2$ (and $\bar{m} = 2$). From Lemma 4.9, $a_3 = 1$. By changing a character χ if necessary we get $\text{wt}(x_3) \equiv 1 \pmod{4}$. Up to permutation $\{1, 2\}$ we also have $\text{wt}(x_1) \equiv 1 \pmod{4}$ and $\text{wt}(x_2) \equiv 3 \pmod{4}$. Then by normalizedness, $a_1 = 1$ and $a_2 = 1$ or 3 . Obviously $\text{wt}(x_4) \equiv 0 \pmod{4}$ and $a_4 = \bar{m} = 2$. Thus a component $C^\sharp(1)$ of $C^\sharp$ is the image of $t \rightarrow (t, t^{a_2}, t, t^2)$, where $a_2 = 1$ or 3 . But if $a_2 = 3$, then $C^\sharp$ can be given by $x_2 - x_1^3 = x_3 - x_1 = x_4^2 - x_1^4 = 0$. Whence an equation of $X^\sharp$ in $\mathbb{C}^4$ in a normalized coordinate system is $\phi = (x_2 - x_1^3)\varphi_1 + (x_3 - x_1)\varphi_2 + (x_4^2 - x_1^4)\varphi_3 = 0$. Since (X, P) is a cyclic quotient, ϕ must contain the term x_4 , a contradiction. Therefore $a_2 = 1$ and $C^\sharp$ is given by $x_2^2 - x_1^2 = x_3 - x_1 = x_4 - x_1x_2 = 0$. By changing coordinates we obtain 4.1.2(iii).

If $d = 4$, then similarly one can see $a_3 = 1$ and $\text{wt}(x_3) \equiv 1 \pmod{8}$. Again an equation of $X^\sharp \subset \mathbb{C}^4$ in a normalized coordinate system contains the term x_4 , so it also contains a monomial $\phi_0 \neq x_4$ such that $\text{wt}(\phi_0) \equiv 0$ and $\text{ord}(\phi_0) = \text{ord}(x_4) = 2$. Then $\phi_0 = x_1x_2$ and $a_1 = a_2 = 1$. Denote $a := \text{wt}(x_1)$. It is easy to see that we can take $a \in \{1, 3\}$. If $a = 1$, then $C^\sharp = \{x_2^4 - x_1^4 = x_3 - x_1 = x_4 - x_1x_2 = 0\}$. If $a = 3$, then one can check that $C^\sharp = \{x_2^2 - x_3^2 = x_2x_3 - x_1^2 = x_4 - x_1x_2 = 0\}$. After changing coordinates we obtain 4.1.3.

To complete the proof we have to show that X can contain at most one Gorenstein singular point if $(S, 0)$ is of type A_1 and contains no Gorenstein singular point if $(S, 0)$ is of type A_3 . Indeed in diagram 3.1 $f^b: (X^b, C^b) \rightarrow (S^b, 0)$ is a Mori conic bundle with reducible central fiber C^b . Therefore by 1.1.1, $\rho(X^b/S^b) > 1$ and each component $C^b(i)$ of C^b generates an extremal ray on X^b over $(S^b, 0)$. Thus each $(X^b, C^b(i))$ is an (not necessary flipping) extremal neighborhood in the sense of [14]. By [14, Theorem (6.2)], we see that $(X^b, C^b(i))$ contains at most one Gorenstein singular point and so does (X, C) because $X^b \rightarrow X$ is étale outside P .

In case 4.1.3 X^b has index 2, whence $(-K_{X^b} \cdot C^b(i)) \in \frac{1}{2}\mathbb{Z}$. On the other hand as in 3.1 $f^b: (X^b, C^b) \rightarrow (S^b, 0)$ is flat. Therefore for a general fiber L of f^b we have $L \equiv r \sum C^b(i)$, where $r \in \mathbb{N}$. Thus $2 = (-K_{X^b} \cdot L) = 4r(-K_{X^b} \cdot C^b(i))$, so $1/2r \in \frac{1}{2}\mathbb{Z}$. It gives us $r = 1$, $L \equiv \sum C^b(i)$. Whence the scheme-theoretical central fiber of f^b over 0 is generically reduced. Then C^b is a complete intersection of Cartier divisors inside X^b . Since C^b has a unique singular point P^b , X^b has exactly one singular point (at P^b) along C^b and so has X (at P). In case (i) of 4.1.2 one can use the same arguments to show that the singular point P is unique. This proves Theorem 4.1.

□

Finally we propose examples of Mori conic bundles over a non-singular base surface (case 4.1.1 of Theorem).

Example 4.18. Let $Y \subset \mathbb{P}_{x,y,z,t}^3 \times \mathbb{C}_{u,v}^2$ is given by the equations

$$\begin{cases} xy - z^2 = ut^2 \\ x^2 = uy^2 + v(z^2 + t^2) \end{cases}$$

It is easy to check that Y is non-singular. The projection $\mathbb{P}^3 \times \mathbb{C}^2 \rightarrow \mathbb{C}^2$ gives us an elliptic fibration $g: Y \rightarrow \mathbb{C}^2$. A general fiber of g is an intersection of two quadrics in $\mathbb{P}^3$ and the central fiber $g^{-1}(0)$ is multiple $\mathbb{P}^1$. Let $\mathbb{Z}_2$ acts on Y by

$$x \rightarrow -x, \quad y \rightarrow -y, \quad z \rightarrow -z, \quad t \rightarrow t, \quad u \rightarrow u, \quad v \rightarrow v.$$

The locus of fixed points consist of the divisor $B := \{t = 0\} \cap Y$ and the isolated point $(x, y, z, t; u, v) = (0, 0, 0, 1; 0, 0)$. Consider the quotient morphism $h: Y \rightarrow X := Y/\mathbb{Z}_2$. By the Hurwitz formula $0 = K_Y = h^*(K_X) + D$, where $h^*(-K_X) = D$. Whence $-K_X$ is ample over $\mathbb{C}^2$. Therefore the quotient $f: X \rightarrow \mathbb{C}^2$ is a Mori conic bundle with irreducible central fiber that contains only one singular point of type $\frac{1}{2}(1, 1, 1)$.

The following example shows that the total space X of a Mori conic bundle can contain two Gorenstein terminal points (cf. [14, 0.4.13.1]).

Example 4.19. Let $Y \subset \mathbb{P}_{x,y,z,t}^3 \times \mathbb{C}_{u,v}^2$ is given by the equations

$$\begin{cases} x^2 = uz^2 + vt^2 \\ y^2 = ut^2 + vz^2 \end{cases}$$

As above the quotient $X := Y/\mathbb{Z}_2(1, 1, 1, 0; 0, 0) \rightarrow \mathbb{C}^2$ is a Mori conic bundle with irreducible central fiber. Singular points of X are one point of type $\frac{1}{2}(1, 1, 1)$ and two ordinary double points.

REFERENCES

- [1] F. Catanese, *Automorphisms of rational double points and moduli spaces of surfaces of general type*, Compositio Math. **61** (1987), no. 1, 81–102.
- [2] H. Clemens, J. Kollár, S. Mori, *Higher dimensional complex geometry*. Astérisque 166 (1988).
- [3] S. Cutkosky, *Elementary contractions of Gorenstein threefolds*, Math. Ann. **280** (1988), 521–525.
- [4] S. Ishii, *On Fano 3-folds with non-rational singularities and two-dimensional base*, Abh. Math. Sem. Univ. Hamburg **64** (1994), 249–277.
- [5] V. A. Iskovskikh, *On the rationality problem for conic bundles*, Mat. Sb. **182** (1991), no.1, 114–121; English transl. Math. USSR-Sb. **72** (1992), no. 1, 105–111.
- [6] V. A. Iskovskikh, *On the rationality criterion for conic bundles*, Mat. Sb. **187** (1996), no. 7, 75–92.
- [7] Y. Kawamata, *Crepant blowing-ups of 3-dimensional canonical singularities and its application to degenerations of surfaces*, Ann. Math. **127** (1988), 93–163.

- [8] Y. Kawamata, K. Matsuda, K. Matsuki, *Introduction to the minimal model program*, in " *Algebraic Geometry, Sendai, 1985*", Adv. Stud. in Pure Math. vol. 10 (1987), pp. 283–360.
- [9] G. Kempf, *Cohomology and convexity*, in " *Toroidal Embeddings I*", Lecture Notes Math. vol. 339 (1973), pp. 41–52.
- [10] J. Kollár, *Standard models of elliptic threefolds*, preprint.
- [11] J. Kollár, Y. Miyaoka, S. Mori, *Rationally connected varieties*, J. Algebraic Geometry **1** (1992), 429–448.
- [12] H. Matsumura, *Commutative ring theory*, Cambridge Univ. Press., Cambridge. 1986.
- [13] S. Mori, *On 3-dimensional terminal singularities*, Nagoya Math. J. **98** (1985), 43–66.
- [14] S. Mori, *Flip theorem and the existence of minimal models for 3-folds*, J. Amer. Math. Soc. **1** (1988), no. 1, 117–253.
- [15] N. Nakayama, *The lower semi-continuity of the plurigeners of complex varieties*, in " *Algebraic Geometry Sendai 1985, T. Oda ed.*", Adv. Stud. in Pure Math., vol. 10, Kinokunya, Tokyo and North-Holland, Amsterdam, 1987, pp. 551–590.
- [16] Yu. G. Prokhorov, *On general elephant problem for three-dimensional $\mathbb{Q}$ -Fano fiber spaces over a surface*, Fundamental'naya i prikladnaya matematika **1** (1995), no. 1, 263–280.
- [17] Yu. G. Prokhorov, *On $\mathbb{Q}$ -Fano fiber spaces with two dimensional base*, preprint MPI 95/33 (1995).
- [18] Yu. G. Prokhorov, *On extremal contractions from threefolds to surfaces*, Warwick preprint 84/1995, (1995).
- [19] Yu. G. Prokhorov, *On general elephant problem for Mori conic bundles*, preprint alg-geom/9608007.
- [20] M. Reid *Minimal models of canonical threefolds*, in " *Algebraic Varieties and Analytic Varieties (S. Iitaka, ed.)*", Adv. Stud. in Pure Math., vol. 1, Kinokunya, Tokyo and North-Holland, Amsterdam, 1983, pp. 131–180.
- [21] M. Reid *Young persons guide to canonical singularities*, in " *Algebraic Geometry, Bowdoin, 1985*, Proc. Symp. Pure Math. vol. 46, 1987, pp. 345–414.

E-mail address: prokhorov@mech.math.msu.su, prokhorov@nw.math.msu.su

ALGEBRA SECTION, DEPARTMENT OF MATHEMATICS, MOSCOW STATE UNIVERSITY,
VOROB'EVY GORY, MOSCOW 117 234, RUSSIA